

## 兔抗 SOCS2 多克隆抗体

中文名称: 兔抗 SOCS2 多克隆抗体

英文名称: Anti-SOCS2 rabbit polyclonal antibody

别名: CIS2; SSI2; Cish2; SSI-2; SOCS-2; STATI2

抗原: SOCS2

储 存: 冷冻(-20℃)

宿 主: Rabbit

相关类别: 一抗

反应种属: Human, Mouse, Rat

标记物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

**Background:** 

This gene encodes a member of the suppressor of cy tokine signaling (SOCS) family. SOCS family members are cytokine-inducible negative regulators of cytokine receptor signaling via the Janus kinase/signal transdu cer and activation of transcription pathway (the JAK/S TAT pathway). SOCS family proteins interact with maj or molecules of signaling complexes to block further signal transduction, in part, by proteasomal depletion of receptors or signal-transducing proteins via ubiquit ination. The expression of this gene can be induced by a subset of cytokines, including erythropoietin, G



	M-CSF, IL10, interferon (IFN)-gamma and by cytokine receptors such as growth horomone receptor.
Applications:	ELISA, IHC
Name of antibody:	SOCS2
Immunogen:	Fusion protein of human SOCS2
Full name:	suppressor of cytokine signaling 2
Synonyms:	CIS2; SSI2; Cish2; SSI-2; SOCS-2; STATI2
SwissProt:	O14508
ELISA Recommended dilution:	1000-2000
IHC positive control:	Human brain
IHC Recommend dilution:	25-100

